Beet Leafhoppers, Heat Injury, and Potential Stink Bug Damage on Seedling Corn

May 22, 2019



In this Issue:

- Beet Leafhoppers: A common threat during the first 6-8 weeks of planting.
- **Heat Injury:** Transplants are susceptible to being burnt on their stems and foliage if they come into contact with black plastic mulch.
- Stink Bugs: Stink bugs can potentially cause damage to young corn stands.

Beet Leafhoppers (Circulifer tenellus)





Leafhopper caught on a yellow sticky trap



Example of row covers to prevent beet leafhoppers

Overview

Beet leafhoppers are prevalent throughout the western United States. Their annual movement follows the wind currents from desert areas in New Mexico and Arizona up into Colorado and Utah. Adults have 3 mm wedge-shaped bodies with wings. Their summer colors can range from a pale green, grey, to tan. During the winter and early spring, they will appear darker in color. In Utah, beet leafhoppers can have 1-2 generations in the spring months (when they are mainly a threat). Then an additional 2-3 generations throughout the summer. Depending on the temperature, females can lay up to 200 eggs. These white/yellow eggs are elongated and slightly curved. Nymphs develop through 5-6 instars, they're spotted with black, red, and brown coloring on their thorax and abdomen.

Damage

Crops are most susceptible in the springtime when the weeds begin to dry up and the pests migrate toward other host crops. Both nymphs and adults have piercing-sucking mouthparts which can causes host plants to shrivel and "burn" under major infestations. Beet Leafhoppers are one of the primary vectors for Curly Top Virus (*Curtovirus*). CTV is unpredictable but widespread across Utah

and is especially threatening to tomato production. Symptoms can include twisted-curled leaves, dark purple veins (not to be confused with phosphorus deficiency), and yellowing of mature leaves. Less common host crops include beets, beans, peppers, cucurbits, and several weed species.

Management

- Use yellow sticky traps to monitor beet leafhopper populations.
- Exclude Beet Leafhoppers (and other potential pests) by using floating row covers.
- Shade your tomato and pepper plants.
- Destroy and remove all plant debris and residue at the end of the growing season.
- "Hide" susceptible plants among resistant plants.
- Beet Leafhopper insecticides registered for <u>commercial</u> use in Utah:

Active Ingredient	Brand Name	Mode of Action	Residual Days	Other Notes
Carbaryl	Carbaryl, Sevin	1A	3	
Dimethoate	Dimate 4E, Dimethoate	1B	6-7	Not for Eggplants
Endosulfan	Thionex ^R	2A		Not for Eggplants
Alpha-cypermethrin	Fastac ^R	3	3	
Bifenthrin + zeta-cypermthrin	Hero ^R , Steed ^R	3	3	
Deltamethrin	Delta Gold ^R	3	5	
Fenpropathrin	Danitol [®]	3	7	
Gamma-cyhalothrin	Declare ^R	3	5	
Lambda-cyhalothrin	Lambda ^R , Paradigm ^R , Silencer ^R	3	5	
Pyrethrins	Pyganic ^{ob}	3		
Zeta-cypermethrin	Mustang ^R	3	7	
Pyrethrin+piperonyl butoxide	Evergren	3	3	
Zeta-cypermethrin + avermectcin BI	Gladiator ^R	3/6	7	
Beta-cyfluthrin + imidacloprid	Leverage 360 ^R	3/4A	7	
Lambda-cyhalothrin + imidacloprid	Kilter [®]	3/4A	5	
Lambda-cyhalothrin +thiamethoxam	Endigo ^R	3/4A	5	
Bifenthrin + abamectin	Athena ^R	3/6	10	
Lambda-cyhalothrin +chlorantraniliprole	Viliam Xpres ^{sR}	3/28	5	
Pyrethrins + azadirachtin	Azera ^{OB}	3/UN	5-7	
Clothianidin	Belay	4A	7	
Imidacloprid	Admire Pro, Couraze, Marathon	4A	5	
Thiamethoxam	Actara, Flagship, Platinum	4A	5	
Spirotetramat	Kontos	23	7	
Chlorantraniliprole+thiamethoxam	Durivo, Voliam Flexi	28/4A	One application per/season	
Beauveria bassiana GHA	Botanigard ^B	NC	2-5	No tomato
Kaolin	Surround ^o	NC	5	
Potassium salts of fatty acids (insecticidal soap	M-Pede	NC	7	
Sodium tetraborohydrate decahydrate	Prev-AM	NC	7-10	
Sucrose octanoate	SucraShield	NC		

• Beet Leafhopper insecticides registered for <u>home</u> use in Utah:

Active Ingredient	Brand Name	Moo Act	le of tion	Residual Days	Other Notes
Malathion	Bonide Malathion, Ortho Max Malathion	1B		5	
Bifenthrin	Binide Eight Flower and Vegetable Granules, Monterey All Natural Mite & Insect Control, Ortho Bug- B-Gon Max	3		7	
Esfenvalerate	Monterey Bug Buster II	3			
Lambda-cyhalothrin	Spectracide Triazicide		3	7	
Permethrin	Bayer Complete Insect Dust, Bonide Eight Vegetable Fruit and Flower	3		3	
Zeta-cypermethrin	GardenTech Sevin	3		14	
Pyrethrins + piperonyl butoxide	Bonide Pyrethrin Garden Insect Pray^B, GardenTech Worry Free Insecticide and Miticide	3		3	
Pyrethrin + canola oil	Monterey Tae Down Garden Spray	3/NC			Only tomatoes
Pyrethrin + sulfure	Bayer Natria Insect, Disease and Mite Control, Bonide Tomato and Vegetable 3 in 1	3/M2		7	Not eggplants
Acetemipirid	Ortho Flower, Fruit, and Vegetable	4A		7	
Capsaicin and related capsaicinoids	Bonide Hot Pepper Wax Insect Repellent	NC			
Oils: canola, clove, cottonseed, garlic, neem, paraffinic, peppermint rosemary	Bayer Natria Multi-insect control ^o , Bonide All Seasons Horticultural and Domant Spray ^o	NC			
Potassium salts of fatty acids (insecticidal soap)	Bayer Natria ^B , Safer Insecticidal Soap ^B , Natural Guard ^B	NC		7	
Additional Resources					

Additional Resources

Beet Leafhoppers (Utah Vegetable Production & Pest Management Guide) Curly Top Virus (Utah Vegetable Production & Pest Management Guide) Curly Top Virus (USU Factsheet) Historical Beet Leafhopper Research in Utah (over 100 years old)

Heat Injury on Transplants



Onions at risk of heat injury laying on black plastic mulch



Transplant at risk of being burned on black plastic mulch



Tomato at risk of being burned on black plastic mulch

Overview

Black plastic mulch is a successful method of weed control and water retention in commercial vegetable production. However, one disadvantage is that it can contribute to heat injury on new transplants. This occurs on clear sunny days that then allow the surface of the black mulch can exceed 130 degrees Fahrenheit.

Damage

When the lower leaves or stem of transplants come into contact with the black plastic mulch at high temperatures this can cause burning to the foliage or stem. This is an abiotic factor and shouldn't be confused with any insect or disease symptoms.

Management

- Depending on the extent of heat injury, replanting might be necessary.
- Reduce risk by improving the contact between the soil and plastic mulch.
- Harden off transplants so they will be more tolerant to heat injury.
- Holes on plastic should be round and even, and transplants should be placed to avoid plant contact with the plastic.

Additional Resources

Use of Plastic Mulch for Vegetable Production (USU Factsheet) Heat Injury with Plastic Mulches (Michigan State Extension) Plasticulture for Commercial Vegetables (North Carolina State Extension)

Potential Stink Bug Damage in Sweet Corn



Stink Bug damage to seedling corn (Kentucky Extension Pest Program)



Brown Marmorated Stink Bug on a corn plant



Stink Bug damage to seedling corn (Kentucky Extension Pest Program)

Overview

Young corn seedlings emerging in the late spring and early summer <u>can be</u> at risk to stink bug damage. Corn species are a part of the wide range of hosts targeted by stink bugs here in Utah. Stink bug species in Utah include the Brown-Marmorated, Brown, Vonchuela, Consperse, Green, One-spotted, Say's, and Uhler. Adults are shield-shaped and lay barrel-shaped egg clusters on host plants leaves. There are typically 2-3 generations within a year.

Damage

In Utah, there has been no reported stink bug damage to seedling corn thus far. However, signs have been reported elsewhere in the country. Knowing the signs can be helpful in assisting to identify stink bugs in the field or garden. Stink bug adults and nymphs have piercing-sucking mouthparts which puncture the stems, leaves, or apical meristems of corn plants. The stink bugs will then inject enzymes into the plant to aid their digestion and fluid removal. Though they can target corn seedlings, stink bugs are more a threat to other host crops during the summer and fall in Utah.

Management

- Attract and conserve natural enemies to stink bugs such as birds, spiders, wheel bugs, assassin bugs, parasitic wasps, and even predatory wheel bugs.
- Monitor for stink bug populations by visual inspection and traps.
- In small sites, hand remove the stinkbugs and discard them into soapy water.
- Remove all weeds that grow near field or garden borders to decrease stink bug attraction.
- Insecticide options are available for stink bugs, but always read the label to ensure they're safe to use on your specific crop.

Additional Resources

Stink Bugs on Corn (Purdue Extension)

Scouting and Managing Stink Bugs in Seedling Corn (North Carolina State Extension) Damage by Stink Bugs Observed in Seedling Corn (Kentucky Extension)